

Information Quality on Developmental Dysplasia of the Hip on Turkish Websites

Türkçe Web Sitelerinde Gelişimsel Kalça Displazisi Hakkındaki Bilgilerin Kalitesi

Hasan Hüseyin Ceylan¹, Barış Çaypınar¹, Mehmet Küçükkoç², Selçuk Uzuner², Fatih Küçükdurmaz³

¹Clinic of Orthopedics and Traumatology, Lütfiye Nuri Burat State Hospital, İstanbul, Turkey ²Department of Pediatrics, Bezmialem Vakıf University School of Medicine, İstanbul, Turkey ³Department of Orthopedics and Traumatology, Bezmialem Vakıf University School of Medicine, İstanbul, Turkey

ABSTRACT

Objective: We aimed to evaluate the reliability of Turkish information achieved from websites about developmental dysplasia of the hip using standardized scientific methods.

Methods: The term "hip dislocation" term was scanned among the most commonly used three internet search engines. The contents of the 10 most frequently clicked sites were copied. These 30 sites were examined. In total, 9 same sites, 3 unrelated sites, and 4 academic sites were eliminated, and 14 sites remained. These sites were scored according to the patient information sheet prepared by the American Academy of Orthopaedic Surgeons. One point was given for every issue included and 0 point was given for no inclusion, over a total of 26 points.

Results: With the search of the term "hip dislocation", 141,000 results on Google, 49,600 on Yahoo, and 46,800 on Bing were found. Results of the content points of the 14 sites that were calculated with the created standard form were determined to be an average of 13.5 points out of 26 (min: 5, max: 22, and SD: 5.02).

Conclusion: We determined that the most visited sites in search engines on hip dislocation contained incomplete information. A variety of applications that aim to increase the reliability of information should be undertaken, or more studies about the development of an accreditation system for Turkish language websites should be performed. (JAREM 2016; 6: 84-7)

Keywords: Hip dislocation, internet, health informatics

ÖΖ

Amaç: Çalışmamızda internette doğumsal kalça çıkığı hakkında sunulan Türkçe bilgilerin güvenilirliğini standardize edilmiş bilimsel yöntemlerle değerlendirmeyi amaçladık.

Yöntemler: "Kalça çıkığı" terimi Türkiye'de en sık kullanılan üç arama motoru olan Google, Yahoo ve Bing üzerinden tarandı. Her arama için arama sonuç listesindeki ilk 10'ar sıradaki site tespit edildi, aynı olan siteler, konuyla ilgisi olmayan siteler ve profesyonellere hitap eden akademik içerikli siteler elendi. Geriye kalan siteler bilgi içeriklerine göre Greene ve ark.nın metoduyla, AAOS'un hazırladığı hasta bilgilendirme formu esas alınarak hazırlanan formdaki alt başlıklarda bilgi içeriğine göre puanlandı. Siteler, hazırlanan standart forma göre içerdikleri her başlık için 1 puan, bahsetmediği konu için de 0 puan verilerek toplamda 26 puanlık bir anket üzerinden değerlendirildi.

Bulgular: "Kalça çıkığı" terimi ile arama yapıldığında Google'da 141.000, Yahoo'da 49.600 ve Bing'de 46.800 sonuç bulundu. Ulaşılan sonuçlar içinde üç ayrı arama motorunda en sık tıklanan ilk 10 site tüm içeriğiyle kopyalandı. Toplamda üç ayrı motorla ulaşılan 30 site- incelemeye alındı. Farklı arama motorlarında çıktığı halde aynı olan (9 site), konuyla ilgisi olmayan (3 site) ve profesyonellere hitap eden akademik içerikli (4 site) olan siteler elendikten sonra geriye 14 site kaldı. Bu 14 sitenin oluşturulan standart formla yapılan içerik puanlaması sonucu sitelerin ortalama puanı 26 üzerinden 13,5 (min: 5, maks: 22, SD: 5,02) olarak belirlendi.

Sonuç: İnternetin gittikçe daha fazla başvurulan bir bilgi kaynağı haline gelmesi, bu kaynaktan edinilen bilginin içeriğinin doğruluğu tartışmasını da beraberinde getirmektedir. Çalışmamızın sonucunda içeriği Türkçe olarak hazırlanmış internet sitelerinde "kalça çıkığı" konusunda arama motorlarında en çok başvurulan sitelerin çoğunluğunun eksik bilgiler içerdiği belirlenmiştir. Bilgi güvenilirliğini arttırmayı hedefleyen çeşitli uygulamalar veya akreditasyon sistemlerinin Türkçe web siteleri için de geliştirilmesi konusunda gerekli çalışmalar yapılmalıdır. (JAREM 2016; 6: 84-7)

Anahtar Kelimeler: Kalça çıkığı, internet, sağlık bilişimi

INTRODUCTION

Internet usage has expanded enormously over the last 10 years in Turkey (Figure 1). Based on findings from the 2014 Turkish Statistical Institute (TurkStat) Internet Use Survey, an estimated 60.2% of households in Turkey had Internet access (1). In 2014, more than half of the population used the Internet for personal purposes from their homes, and 53.8% of Turkish adults used the Internet to access information. According to data for 2009, nearly 45%

This study was presented at the İstanbul Orthopaedics Meeting, 24-27 May 2013, İstanbul, Turkey. Bu çalışma İstanbul Ortopedi Buluşması'nda sunulmuştur, 24-27 Nisan 2013, İstanbul, Türkiye.

of people accessed the Internet for health-related information (2). Internet users went online to search for health-related information as other subjects that they want to take information. The population in Turkey is younger and also prolific (1).

The Internet is a worldwide data-sharing platform that has both advantages and limitations when used as a source of information (3). The Internet is an easy, quick, and inexpensive platform for collecting information on any issue. Many types of available

Address for Correspondence / Yazışma Adresi: Dr. Hasan Hüseyin Ceylan, E-mail: drhhc@yahoo.com

Received Date / Geliş Tarihi: 26.06.2015 Accepted Date / Kabul Tarihi: 29.07.2015 © Copyright 2016 by Gaziosmanpaşa Taksim Training and Research Hospital. Available on-line at www.jarem.org © Telif Hakki 2016 Gaziosmanpaşa Taksim Eğitim ve Araştırma Hastanesi. Makale metnine www.jarem.org web sayfasından ulaşılabilir. DDI: 10.5152/jarem.2015.827

data exist on the Internet for users (4), allowing patients to obtain information about diseases and therapies themselves. Sometimes, this might include important aspects of self-diagnosis and self-management. Patients can also learn about the emerging aspects of the diagnosis of common health problems widely observed in our population.

Because of high birthrates and cultural preferences, such as swaddling, developmental dysplasia of the hip (DDH) is an important problem in Turkey. Early diagnosis, particularly in the first three weeks after birth, improves the outcomes in DDH. During this short period, nonsurgical treatment can be used to rectify the hip joint (5). Therefore, the quality of available information is extremely important for the early diagnosis and optimal treatment of DDH. Similar to the universal trend, the number of people in Turkey obtaining information about developmental diseases such as DDH from the Internet has been increasing. In this study, we aimed to investigate the quality of information about DDH on Turkish websites that may affect the insight of the patient's relatives.

METHODS

A simple Internet search using the search term "kalça çıkığı" (Turkish for DDH) was performed on March 15, 2014 using the three most popular search engines: Google (http://www.google.com), Yahoo (http://www.yahoo.com), and MSN (http://www.bing.com), which account for more than 95% of Internet searches (6). The first 10 websites from each search engine were identified based on the researched fact that people generally view only the first 10 results from Internet searches (Figure 2) (7). Thus, 30 websites were identified.

Websites were classified into the following three categories: public (easy for any person to understand), academic (addressed to health professionals), and sites with no focus on DDH. We only evaluated the public sites because the academic and unrelated sites did not address the information needs of the general public about DDH. For each included site, relevant data were scored according to the ingredient questionnaire previously described by Greene et al. (8) (Table 1). The sum of each included parameter represented the average score for that site.

Consecutive data analysis was performed in a number of stages. First, two qualified pediatric orthopedic surgeons generated a content quality score according to the DDH Patient Educating Form of the American Academy of Orthopedic Surgeons (AAOS) (9). All search data and details were saved. Then, two other orthopedic surgeons evaluated the information and graded it according to the criteria based on the AAOS DDH Patient Educating Form. The DDH questionnaire includes basic information needed by patients in a format they can understand. All steps of our study was designed according to Declaration of Helsinki.

RESULTS

For the search term kalça çıkığı (i.e., hip dislocation or DDH), we obtained approximately 141,000 results from Google, 49,600 from Yahoo, and 46,800 from MSN. The 10 most frequently clicked sites per search engine were copied with all their content. Thus, 30 sites were accessed for further examination. After eliminating duplicate sites (n=9), unrelated sites (n=3), and sites with aca-

demic content more suited to professionals (n=4), 14 sites were identified for further analysis. Using the standard form, the average content score for the 14 sites was calculated as 13.5 out of 26 points (range, 5–22; SD, \pm 5.02). The site www.doktorum.com had the highest score in this study. Other top ranking sites were www. wikipedia.com and www.ortopedikbilgi.com. Only five websites included basic information about the disease. Commercial sites had significantly more accessibility violations.

DISCUSSION

Hip dysplasia has been defined as the lack of correct orientation of the femoral head and acetabulum. DDH is one of the most widely discussed abnormalities in neonates and has a multifactorial etiology. The reported incidence of DDH varies, but is reported at 1.5–20 cases per 1000 births, possibly because of the variability of diagnostic methods, tool quality, and timing (10). Many factors, including genetics, intrauterine position, breech presentation, sex, race, family history, associated skeletal abnormalities, first-born, hormonal factors, environmental factors, and oligohydramnios, are known risk factors for DDH (11, 12). Family history and inheritance are also important, and several postnatal factors play important roles in the etiology of DDH. For example, swaddling forces the hips into adduction and extension, which is unfavorable for the developing hip joint. This postnatal practice is common in most of Northeastern Anatolia. In Anatolia, the incidence of swaddling is nearly 47%, and the incidence of DDH is estimated to be nearly 1.4% in Turkey (13).

The Internet has been an important information source on various issues, including medical or health issues, surgical therapy,







drug therapy, healthcare technology, and alternative paramedical therapy. Many parents look for information about DDH from various sources. Increased use of the Internet aids the families of patients with DDH to obtain information from the web. However, there is a lack of content control in web sources, and families sometimes obtain incorrect information. Moreover, it has been reported that 85% of parents do not know the full details of the disease and believe that DDH can be treated after their child has learned to walk (14, 15).

Health information accessed from the Internet is important in

17). Berland et al. (18) claimed that accessing health information from search engines is not a suitable method, and recommended systematic evaluation and control of websites concerning health issues. Physicians should be aware of the quality and accuracy of information available on the Internet as this can impact medical decision-making and healthcare decisions in families.

In general, patients use popular search engines (e.g., Google) rather than professional search engines (e.g., PubMed) for health-related literature (18, 19), and rarely read the results after the first 10 ranked websites. Therefore, decisions are made on the basis of these ranked sites (7). The quality of healthcare-related infor-

Table 1. Questionnaire interrogating information details about DDH Score Headings of information **Subheadings** 1 Description Normal anatomy of the hip Developmental dysplasia (dislocation) of the hip (DDH), i.e., 1 the hip joint has not formed normally At birth or soon after can occur increase in activity Diagnosis is made by examination by the general pediatrician Causes Girls, first-born children, left hips, and breech presentation are associated with increased risk Family history of DDH (parents or siblings) and oligohydramnios (low levels of amniotic fluid) Symptoms Some babies born with a dislocated hip will show no outward signs Legs of different lengths Limping, walking on the toes, waddling, or duck-like gait Uneven skin folds Less mobility or flexibility on one side Examination Clinical examination is performed first Infants monitored using USG After 6 months, monitor using X-ray Nonsurgical treatment Bandage treatment of newborn infants 1-6 months' bandaging or individual plaster treatment Usually performed after 6 months Surgical treatment There are options for intervention by age Treatment is performed with anesthesia Open surgery may be necessary After the intervention, a plaster cast is made Controls are monitored by X-ray Complications Calcification can occur despite treatment There can be a delay in walking after treatment The Pavlik harness and other positioning devices may cause skin irritation around the straps Difference in leg length may remain TOTAL 26 DDH: developmental dysplasia of the hip; USG: ultrasonography

helping patients make decisions and affects their preferences (16, of the second

mation on the Internet is presently uncontrolled, and warnings regarding this have been issued (20). Healthcare resources in Turkish websites need to be evaluated and accredited by an independent institutional body for patients' benefits, similar to the Health on Net Foundation (HON) panel that independently evaluates the quality of medical websites with specific guidance (21).

CONCLUSION

Independent, external validation and accreditation of websites that provide healthcare information is mandatory to optimize guidance to the Turkish population. Information on the Internet must be reinforced by medical professionals educating the population, thereby preventing incorrect practices, irreversible prejudices, and limiting the effects of commercial goals.

Ethics Committee Approval: Authors declared that the research was conducted according to the principles of the World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects", (amended in October 2013).

Informed Consent: No patient data used fort his study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - H.H.C., B.Ç., M.K., S.U., F.K.; Design -H.H.C., B.Ç., M.K., S.U., F.K.; Supervision - H.H.C., B.Ç., M.K., S.U., F.K.; Resources - H.H.C., F.K., M.K.; Materials - H.H.C., M.K., S.U.; Data Collection and/or Processing - H.H.C., S.U., M.K., F.K.; Analysis and/or Interpretation - H.H.C., F.K., B.Ç.; Literature Search - H.H.C., M.K., B.Ç.; Writing Manuscript - H.H.C., S.U., B.Ç.; Critical Review - H.H.C., F.K., B.Ç.

Acknowledgements: We thank to Talat Bulut for English grammar edition of this paper.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Yazarlar çalışmanın World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects", (amended in October 2013) prensiplerine uygun olarak yapıldığını beyan etmişlerdir.

Hasta Onamı: Çalışmamızda hasta verisi kullanılmamıştır.

Hakem Değerlendirmesi: Dış bağımsız.

Yazar Katkıları: Fikir - H.H.C., B.Ç., M.K., S.U., F.K.; Tasarım - H.H.C., B.Ç., M.K., S.U., F.K.; Denetleme - H.H.C., B.Ç., M.K., S.U., F.K.; Kaynaklar -H.H.C., F.K., M.K.; Malzemeler - H.H.C., M.K., S.U.; Veri Toplanması ve/ veya İşlemesi - H.H.C., S.U., M.K., F.K.; Analiz ve/veya Yorum - H.H.C., F.K., B.Ç.; Literatür Taraması - H.H.C., M.K., B.Ç.; Yazıyı Yazan - H.H.C., S.U., B.Ç.; Eleştirel İnceleme - H.H.C., F.K., B.Ç.

Teşekkür: Metnin İngilizce gramer edisyonu için yardımcı olan Talat Bulut'a teşekkür ederiz.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

REFERENCES

- TurkStat (TUIK) annuals. Available from: http://www.tuik.gov.tr/Pre-HaberBultenleri.do?id=16198 Date last updated: August 22 2014. Date last accessed: April 23 2015.
- Son üç ay içinde internet kullanan bireylerin interneti kişisel kullanma amaçları verisi. Bilim, Teknoloji ve Bilişim Ocak-Mart 2009 Dönemi verileri istatistikleri. Türkiye İstatistik Yıllığı, p.428-42.
- Seale C, Charteris-Black J, MacFarlane A, McPherson A. Interviews and internet forums: a comparison of two sources of qualitative data. Qual Health Res 2010; 20: 595-606. [CrossRef]
- Shah SG, Robinson I. Patients' perspectives on self-testing of oral anticoagulation therapy: content analysis of patients' internet blogs. BMC Health Serv Res 2011; 3: 11-25. [CrossRef]
- Dezateux C, Godward S. Evaluating the national screening programme for congenital dislocation of the hip. J Med Screen 1995; 2: 200-6.
- 2010 Search engines statistics that people commonly visits. Also available at http://www.seoconsultants.com/search-engines/ Date last updated: August 9 2010. Date last accessed: April 23 2015.
- Silverstein C, Marais H, Henzinger M, Moricz M. Analysis of a very largeweb search engine query log. ACM SIGIR Forum 1999; 33: 6-12. [CrossRef]
- Greene DL, Appel AJ, Reinert SE, Palumbo MA. Lumbar disc herniation: evaluation of information on the internet. Spine (Phila Pa 1976) 2005; 30: 826-9. [CrossRef]
- AAOS DDH info for patients http://orthoinfo.aaos.org/topic. cfm?topic=A00347 Date last updated: October 2013. Date last accessed: April 23 2015.
- Bialik V, Bialik GM, Blazer S, Sujov P, Wiener F, Berant M. Developmental dysplasia of the hip: a new approach to incidence. Pediatrics 1999; 103: 93-9. [CrossRef]
- Peled E, Eidelman M, Katzman A, Bialik V. Neonatal incidence of hip dysplasia ten years of experience. Clin Orthop Relat Res 2008; 466: 771-5. [CrossRef]
- 12. Storer SK, Skaggs DL. Developmental dysplasia of the hip. Am Fam Physician 2006; 74: 1310-6.
- Bayındır Ş, Tanış Z. Boş batın filimlerinde tesadüfen karşılaşılan doğuştan kalça çıkığı ve diğer patolojileri. Hacettepe Tıp Cerrahi Bülteni 1970; 3: 220-31.
- Bursalı A, Bulut A, Gökçay G. Gelişimsel kalça displazisinde olası bir risk faktörü: bebeği baş aşağı durulama. Çocuk Dergisi 2006; 6: 53-61.
- 15. Bursalı A, Bulut A, Gökçay G. Gelişimsel kalça displazisi ve kadınların bu konudaki bilgi düzeyi. Çocuk Dergisi 2006; 6: 172-8.
- Basch EM, Thaler HT, Shi W, Yakren S, Schrag D. Use of information resourcesby patients with cancer and their companions. Cancer 2004; 100: 2476-83. [CrossRef]
- Black PC, Penson DF. Prostate cancer on the Internet-information or misinformation? J Urol 2006; 175: 1836-42. [CrossRef]
- Berland GK, Elliott MN, Morales LS, Algazy JI, Kravitz RL, Broder MS, et al. Health information on the Internet accessibility, quality, and readability in English and Spanish. JAMA 2001; 285: 2612-21. [CrossRef]
- 19. Krempec J, Hall J, Biermann JS. Internet use by patients in orthopaedic surgery. Iowa Orthop J 2003; 23: 80-2.
- Crocco AG, Villasis-Keever M, Jadad AR. Two wrongs don't make a right: harm aggravated by inaccurate information on the Internet. Pediatrics 2002; 109: 522-3. [CrossRef]
- Health on the Net Foundation. HON code of conduct (HON code) for medical and health websites. www.hon.ch/HONcode Date last updated: August 25 2014. Date last accessed: April 23 2015.