

Outcomes of Post-traumatic Stiff Elbow Arthrolysis

Erdem Özden¹ , Kahraman Öztürk² , Bilal Birkan Demir² , Ahmet Utku Gürün⁴ , Murat Demiroğlu³ , Serkan Aykut² 

¹University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital, Clinic of Orthopedics and Traumatology, İstanbul, Turkey

²İstanbul Baltalimanı Metin Sabancı Bone Diseases Training and Research Hospital, Clinic of Hand and Upper Extremity Surgery, İstanbul, Turkey

³İstanbul Medeniyet University Göztepe Training and Research Hospital, Clinic of Orthopedics and Traumatology, İstanbul, Turkey

⁴İstanbul Bağcılar Training and Research Hospital, Clinic of Orthopedics and Traumatology, İstanbul, Turkey

Cite this article as: Özden E, Öztürk K, Demir BB, Gürün AU, Demiroğlu M, Aykut S. Outcomes of Post-traumatic Stiff Elbow Arthrolysis. JAREM 2020;10(1): 75-81

ABSTRACT

Objective: In this study, the results of open release surgeries performed for the stiff elbows which were caused by intrinsic, extrinsic or by combination of both factors were evaluated.

Methods: Twenty-three elbows of 22 patients who could not perform functional elbow motions were surgically treated between January 2005 and December 2012. The mean age was 30.6±11.4 years (16 to 67 years), and the average follow-up period was 81 months (58 months to 12 years). Elbow arc of motions was recorded pre- and postoperatively. Patients were evaluated clinically by using quick Disabilities of the Arm, Shoulder and Hand (DASH) self-report questionnaire, and MAYO elbow performance scores.

Results: The mean preoperative elbow arc of motion was 42 degrees (0-90), and it became 109 degrees (70-140) postoperatively after at least 58-month follow-up. The average increase was 67.8±25 degrees (30-125). The mean Quick DASH score was 18.2±12.7 (6.8 to 63.6). According to MAYO elbow rating system, one patient who had experienced infection following the initial fracture treatment had poor result, three patients had fair results, 17 patients had good results, and one patient who had bilateral elbow stiffness had excellent results.

Conclusion: Although good results can be achieved by open release of the stiff elbows, one must keep in mind that preventing the stiffness rather than solving this problem would bring better results. As 15 patients with stiff elbows in this study had a previous surgery performed for an elbow region fracture, stabile fixation, meticulous hemostasis, shorter immobilization period and early rehabilitation can be considered as the most important steps for avoiding elbow stiffness where surgery is indicated.

Keywords: Elbow, stiff, arthrolysis, post-traumatic

INTRODUCTION

The elbow joint is very congruent with three articulations: the ulnohumeral joint, the radiocapitellar joint, and the proximal radioulnar joint. The stability of the elbow is provided by static and dynamic stabilizers. The ulnohumeral articulation, the anterior bundle of the medial collateral ligament, and the lateral collateral

ligament complex are acting as the primary static constraints, and the radiocapitellar articulation, the common flexor tendon, the common extensor tendon, and the capsule are acting as secondary static constraints. The dynamic stabilizers are the muscles crossing the elbow (1-3).

Morrey et al. (1) pointed out that 30 to 130 degrees of flexion with 50 degrees pronation and 50 degrees supination may provide

ORCID IDs of the authors: E.Ö. 0000-0003-0765-5635; K.Ö. 0000-0002-7644-659X; B.B.D. 0000-0003-3449-3870; A.U.G. 0000-0001-5326-7161; M.D. 0000-0001-7992-2662; S.A. 0000-0003-1028-2120.



Corresponding Author: Erdem Özden

E-mail: ozdenerdem@hotmail.com



Received Date: 21.01.2019 **Accepted Date:** 03.04.2019

©Copyright 2020 by University of Health Sciences Turkey Gaziosmanpaşa Training and Research Hospital. Available on-line at www.jarem.org

a functional range of motion (ROM) for most of the daily living activities. We may call the elbow "stiff" if the hand cannot be used functionally due to the loss of elbow motion and this functionality differs for each individual.

The etiology of elbow stiffness must be well understood while planning the treatment. Anatomically the reasons can be classified as extrinsic, intrinsic or combination of both. The intrinsic contracture term is used to determine that the pathology involves the articular surface such as articular adhesions, arthritis, osteophytes, loose bodies, malunions of intra-articular fractures and proliferative synovitis. The extrinsic contracture term defines the periarticular pathologies such as capsule, ligaments, muscles and skin contractures surrounding the joint and also the heterotopic bone formation within these structures. The most common type of elbow contracture is the post-traumatic stiff elbow which usually involves both of the intrinsic and extrinsic pathologies (4,5).

The conservative treatment of stiff elbow includes therapeutic heat application, myofascial soft tissue mobilization, progressive ROM exercises and contracture splinting. Surgery is considered when conservative treatments fail. Surgical treatment options have a spectrum from arthroscopic or open anterior and posterior capsular release to total elbow arthroplasty (6,7). Molecular pathogenesis of the stiffness is still under investigation and immunohistochemical studies warrants further investigation (8,9).

The purpose of this study is to report the functional outcome of the surgically treated stiff elbows.

METHODS

We reviewed 23 stiff elbows of 22 patients who underwent surgical release between January 2005 and December 2012. Conservative treatment methods were tried for at least six months for all of the patients. Surgery was considered for the patients who could not perform functional elbow motions and did not respond to conservative treatment methods. Plain radiographs were obtained to assess the heterotopic bone formations, articular surfaces, hardware positions and the healing of initial fractures. Computerized tomography (CT) scans were also obtained before the treatment to evaluate the articular surfaces more accurately.

The mean age of the patients was 30.6 ± 11.4 years (16 to 67 years), six of the patients were female and 16 were male. The average follow-up period was 81 months (58 months to 12 years). Elbow arc of motions was recorded preoperatively and postoperatively. Patients were evaluated clinically by using quick Disabilities of the Arm, Shoulder and Hand (DASH) self-report questionnaire, and MAYO elbow performance scores.

Only post-traumatic stiff elbows were included in this study; nine patients experienced radial head fractures, four patients had complex elbow fractures (humerus lower end fracture dislocation accompanying ulna or radius fractures), four patients had humerus lower end fractures, three elbows of two patients had heterotopic

ossification (HO) due to head and elbow trauma, one patient had olecranon fracture, one patient had radial head chronic luxation, and one patient had radial head fracture with humero-ulnar luxation. Fifteen patients with fractures underwent surgical intervention after the fractures occurred. The mean time elapsed from the initial trauma to surgical release was 21.4 months. No joint instability was observed before and after the treatment. The etiologies of the stiffness were also demonstrated in Table 1.

All surgical arthrolyses were performed by one of the senior hand and upper extremity surgeon at the author's clinic.

All surgeries were performed under regional block anesthesia and indwelling brachial plexus catheters were placed for postoperative pain management. Lateral incisions alone were used in eleven patients, medial incisions alone in four patients, single anterior incisions in three elbows of two patients, medial and lateral incisions were combined in four patients, and a single posterior incision was used for one patient. Anterior capsulectomy (Figure 1) was performed for extension deficit and posterior capsulectomy (Figure 2) was performed to gain flexion (Figure 3, 4). Additionally, radial head excision was also performed for four patients, HO excision for three elbows of two patients, radial head prosthesis excision for three patients, nervus ulnaris anterior transposition (Figure 1) for four patients and nervus ulnaris release for five patients. After soft tissue and heterotopic bone excisions, olecranon fossa and the coronoid fossa were explored to check if any osteophytes or callus formation making a mechanical block remained. If the extension block was due to posterior impingement, widening the olecranon fossa by the help of a high-speed burr and also the excision of the tip of the olecranon while preserving the triceps tendon insertion were generally needed. If there was any hardware to be removed, the removal was performed at the end of the operations. The tourniquets were released, surgical hemostasis was achieved by cauterization. After applying closed suction drainage, skin closures were performed. The degree of flexion/extension and also supination/pronation achieved under anesthesia were measured and noted. The elbow was usually splinted in maximal extension, but if the patient was suffering only for lack of flexion and already had full extension before the operative release, the elbow was splinted in maximal flexion for 24 hours. Progressive passive ROM exercises was started in the following day and the drains were left in place until the drainage stopped, which was usually observed in the third day after surgery. Patients received a therapy program immediately after discharge.

Patients' data were collected with permission, by reviewing the medical records. This study was conducted in accordance with the Declaration of Helsinki. Ethics committee approval was received for this study from the ethics committee of University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital (approval number: 125; date: 16.01.2019). Informed consent was obtained.

Table 1. Patient's descriptive data

Patient No	Age, y Sex side	Pre-op flex/ ext sup/pron	Post-op final flex/ ext sup/ pron	Gained arc of motion (flex/ext)	Approach	Etiology	Pre-op. ulnar neuropathy	DASH score pre-op	DASH scores post-op	MAYO scores Pre-op	MAYO scores post-op
1	40/M/right	140/-50	150/-10	50	Lateral	Radial head fracture	-	50	20.50	70	85
Left elbow of 2	16/M	90/-40	130/0	80	Anterior	Myositis ossificans	-	56.8	27.3	60	100
Right elbow of 2	-	60/no motion (ankylosis)	130/-10	120	Anterior	Myositis ossificans	-	61.4	6.8	50	100
3	27/F/left	80 / -30	130/-10	70	Medial	Radial head fracture	+	54.5	20.50	55	85
4	25/M/left	90 / -40	110/-25	35	Lateral	Radial head + capitellum fracture	-	61.4	34.10	50	65
5	35/M/right	90 / -45	110/0	65	Lateral + Medial	Complex elbow fracture	+	65.9	15.90	50	85
6	44/M/left	90 / -40	130/-20	60	Medial	Humerus lower end fracture	-	63.6	13.6	50	85
7	35/M/left	60 / -40	80 / -10	50	Lateral	Radial head & coronoid fracture	-	81.8	63.6	30	50
8	16/M/right	120/-55	130 / -10	55	Lateral + Medial	Humerus lower end fracture	-	68.2	6.8	65	85
9	36/M/right	110/-40	135 / -10	55	Medial	Olecranon fracture	-	65.9	22.7	70	85
10	29/M/right	70/-40	130/-10	90	Lateral	Radial head fracture	-	70.5	11.4	50	85
11	26/M/left	100/-60	100/-30	30	Lateral	Complex elbow fracture	-	77.3	31.8	40	60
12	32/F/right	90/-70	120/-40	60	Posterior	Complex elbow fracture	-	77.3	29.50	40	60
13	24/M/right	120/-45	135/-20	40	Lateral + Medial	Radial head fracture	-	63.6	11.4	55	85
14	20/M/left	80/-45	120/-20	65	Lateral	Humerus lower end fracture	-	70.5	13.6	50	85
15	47/F/right	110/-50	130/-20	50	Lateral	Radial head fracture	-	63.6	11.4	50	85
16	24/F/right	Fixed at 90 degree flexion	135/-20	115	Anterior	Myositis ossificans	-	77.3	13.6	45	85
17	32/F/right	Fixed at 90 degree flexion	135/-10	125	Lateral + Medial	Humerus lower end fracture	+	79.5	11.4	45	85
18	67/M/right	90/-45	140/-20	75	Medial	Radial head fracture + humero-ulnar luxation	+	81.8	9.1	45	85
19	28/M/left	90/-60	120/-30	60	Lateral	Radial head fracture	-	79.5	13.6	45	80
20	31/F/right	80/-40	130/-20	70	Lateral	Radial head chronic luxation	-	77.3	15.9	50	80
21	30/M/right	100/-50	140/-20	70	Lateral	Radial head fracture	-	50	6.8	50	85
22	24/M/left	110/-50	130/-20	70	Lateral	Complex elbow fracture	-	56.8	9.1	55	85

DASH: Disabilities of the arm, shoulder and hand, M: male, F: female

Statistical Analysis

We have compared the preoperative and post-operative flexion and extension degrees, DASH scores, and the MAYO elbow performance scores with the Wilcoxon rank-sum test by using SPSS 22.0 (SPSS Inc, IBM, Chicago, IL) software. P-values less than 0.05 were regarded as statistically significant.

RESULTS

The mean preoperative elbow flexion was 93.4 ± 19 degrees (ranges from 60 to 140 degrees) and lack of extension was 51 ± 15.2 degrees (30 to 90 degrees). The average postoperative flexion of elbow was 126 ± 14.8 degrees (80 to 150 degrees) and lack of extension was 17 ± 9.4 degrees (0 to 40 degrees) after a mean follow-up period of 81 months. Preoperative mean arc of motion was 42 degrees (0 to 90 degrees), and it became 109.3 degrees (70 to 140 degrees) postoperatively after at least 58-month follow-up. The average increase was 67.8 ± 24.9 degrees (30-125 degrees). The preoperative mean Quick DASH score was 69 ± 9 points, and it was calculated as 18.2 ± 12.7 (6.8 to 63.6) at the last visit. These changes were found statistically significant. According to MAYO elbow rating system, one patient who had experienced infection following the initial surgery for the fracture treatment had poor results, three patients had fair results, 17 patients had good results, and one patient had excellent results for both of his elbows. Ulnar nerve related symptoms of four patients were recovered after anterior transposition of the nerves. No infection, instability or recurrence of HO was observed after the releases.

DISCUSSION

In this study, we described our experience in surgical release of stiff elbows. Before deciding surgical intervention to release the contracture, the pathology and also the expectations of the patient must be well understood and the patient must be aware of the risks of the procedure such as pain, instability, weakness and worsening of the ulnar nerve related symptoms.



Figure 1. Posterior capsular excision by using the "medial over the top" technique



Figure 2. Anterior capsular excision




Figure 3. Excised posterior capsule




Figure 4. Excised anterior capsule

Preoperative assessments about the articular cartilage and the congruity of the joint would change the operative procedure from basic capsular release to more complex interventions such as fascial graft interposition. Therefore, CT of the joint must be obtained before making the decision. In our cases, no fascial graft interposition was needed, CT scans were also obtained before the treatment.

Pain would seldom be present with a stiff elbow and if present, it is generally seen at the end of flexion or extension; however, if pain is elicited during the whole arc, it should be associated with joint incongruity, synovitis, loose bodies, and arthrosis (10), and this finding could change the type of operation to be performed. In our study, there was no motion at three elbows of two patients who had HO. And the remaining patients were feeling pain only at the end of the arc of motion.

There are several operative techniques determined for the arthrolysis of the contracted elbows and the surgeon dealing with stiff elbows must be familiar with different types of approaches as one incision should not be enough for an adequate release in some cases (11). Manipulation under anesthesia, elbow arthroscopy, synovectomy, excision of radial head, triceps lengthening, excision of ectopic bones, anterior and posterior capsulectomy, distraction arthroplasty, fascial interposition arthroplasty and total elbow arthroplasty are the described procedures for stiff elbow management (11-13).

Although arthroscopic release of the stiff elbows was reported as a safe and effective method by several authors, the indications are limited and cannot be used for the elbows where the anatomy is altered (14-16). We have not performed any arthroscopic release for the patients who were included in this study.

Operative approaches were varying depending on the preoperative and intraoperative findings. Previous incisions, location of the HO and presence of ulnar neuropathy were some of the factors affecting the decision making about the appropriate approach.

For most of our cases, the lateral column procedure was performed, in which a limited lateral approach was used (4). Medial "over-the-top" approach, which was popularized by Mansat et al. (17), was our preferred approach if ulnar nerve symptoms were present pre-operatively and anterior subcutaneous transpositions of the ulnar nerves were performed in those cases. We prefer release and anterior transposition of the ulnar nerve in patients who have preoperative nerve related symptoms, whose elbows are contracted in an extension position for a long time or the amount of postoperatively expected flexion gain is significant.

Another method is a transhumeral, so-called Outerbridge-Kashiwagi (O-K) method, which was originally described by Kashiwagi in 1978. In this technique, a posterior incision is used and after posterior capsulectomy, the olecranon fossa is fenestrated, then anterior capsulectomy can be performed through this hole (18). Morrey (19) have modified this technique and named it as ulnohumeral arthroplasty (UHA) and used a

triceps elevation technique rather than splitting it. Hertel et al. (20) have described a sequential arthrolysis which starts with the standard O-K method and continued with a limited lateral and a limited medial approach if needed for adequate release. Posterior incisions or anterior incisions to address the heterotopic bones can also be used. In our experience, we have found lateral or medial approach useful if the olecranon and its fossa have to be reshaped due to osteophyte formation. Neither O-K method nor UHA was needed in our cases.

For long-standing contractures of the elbow, muscle tightness would also restrict motion and Bhattacharya have suggested triceps and brachialis muscle mobilization as a solution, but this will result in loss of strength and will risk the necessary post-operative rehabilitation program and must be avoided if possible (12). It is also important to preserve the primary constraints of the elbow to avoid laxity and the need for an external fixator device. We did not apply any external fixator to our patients.

If severe arthrosis is the cause of stiffness, releasing the soft tissues would improve the ROM but unfortunately the pain would continue being the most important complaint of the patient and he or she would seek the preoperative condition of the elbow. Therefore, it is important to realize this condition and perform other types of operations like arthrodesis, distraction with fascial interposition arthroplasty or total elbow arthroplasty, which should only be used in some selected patients (13,21,22). Our patient population did not have severe arthrosis; therefore, contracture releases were found to be enough.

Lindenhovius et al. (23) stated that the release in elbows with HO would have better results than the stiff elbows without HO. Our results with HO resections support this statement. In case of total bony ankylosis, the muscle control and strength must be carefully evaluated as these patients would lose the gained arc if they cannot actively flex or extend their elbows after releasing the stiffness.

In a report by Liu et al. (24), the mean improvement in total elbow flexion/extension motions was 80 degrees documented in 11 patients by using combination of lateral and medial approaches and hinged external fixation. In another study performed by Ehsan et al. (25), he documents a total improvement of 58 degrees of elbow flexion/extension motions. Of interest, 68 of 77 patients demonstrated radiographic evidence of HO and 53 patients (69%) achieved a total arc of motion ≥ 100 degrees. Another report by Kayalar et al. (26) represents an increase of 66 degrees in total flexion/extension arc. The average increase in our study was 67.8 degrees (30-125 degrees), which is comparable with others.

The post-operative physical therapy program of these patients is of paramount importance because maintenance of the gained ROM will affect the overall results. Unfortunately, there is no consensus about the most effective therapy program in the literature (6). Some authors have used immediate postoperative continuous passive motion (CPM), interscalene blocks and corrective splinting modalities but some authors did not ever

use the CPM (5,15,23,24). In our clinic, we use static splinting for the first post-operative day in an elevated position to decrease the soft tissue swelling, and the position of the elbow in the long arm splint depends on the patient's preoperative findings. If the operation was performed for a flexion contracture with or without lack of flexion, the elbow was splinted in maximal extension position that was gained in the operation room. If the aim was to gain only flexion, the position of the splint was set in flexion. We believe that educating the patient about passive exercises is a very effective method if he/she is compatible and capable of using his/her other upper extremity. After 24 hours, the patient begins passively flexing the elbow to the end point and stands in that position for fifteen minutes. After muscle relaxation, the patient can flex the elbow a little bit more and holds that position for another 15 minutes. Then does the same for extension exercises and wears the splint when got tired. The splints are worn every resting time and after three weeks worn only nightly. Indwelling brachial plexus catheter was used for pain management during the therapy program for two days and followed by non-steroidal anti-inflammatory agents. Indomethacin 25 mg three times daily was preferred if there were ectopic bones.

Charalambous and Morrey (8) emphasized the importance of molecular pathogenesis of the stiffness in a review of posttraumatic elbow stiffness. In a study conducted by Hildebrand et al. (27), anterior capsules of contracted elbows and healthy elbows of organ donors were compared and immunohistochemical study revealed that myofibroblast numbers were significantly elevated in the contracted elbow capsules. Cohen et al. (28) also investigated the contracted capsules and compared to normal capsules of donors and demonstrated that the contracted capsules were thicker and cytokine levels (MMP-1, MMP-2 and MMP-3) were significantly higher than normal capsules, while collagen type III had decreased levels. According to these findings, they stated that the contracture tissue formation mechanism was different than normal wound healing. Hildebrand et al. (9) also assessed joint myofibroblasts, nerve fibers containing neuropeptides and mast cells by immunohistochemistry in another study and found significantly greater expression in the contracture capsules. They suggested that a manipulation to myofibroblast-mast cell-neuropeptide fibrosis link should be a solution which warrants further investigation.

Molecular pathogenesis of the stiffness is still under investigation and immunohistochemical studies warrants further investigation.

CONCLUSION

Although good results can be achieved by open release of the stiff elbows, one must keep in mind that preventing the stiffness rather than solving this problem would bring better results. As 15 patients with stiff elbows among the 23 patients included in our study had a previous surgery performed for an elbow region fracture, we may conclude that stable fixation, meticulous hemostasis, minimal immobilization and early rehabilitation must

be achieved for avoiding elbow stiffness where surgical treatment is indicated.

However, any patient with elbow trauma must be informed about the possibility of stiffness. And, we believe that the basic scientific investigations should focus on the prophylaxis of joint stiffness in the future.

Ethics Committee Approval: Ethics committee approval was received for this study from the ethics committee of University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital (approval number: 125; date: 16.01.2019).

Informed Consent: It was obtained.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - E.Ö., K.Ö., S.A.; Design - B.B.D., M.D., A.U.G.; Supervision - B.B.D., M.D., A.U.G.; Resources - B.B.D., M.D., A.U.G.; Materials - K.Ö.; Data Collection or Processing - E.Ö., K.Ö., S.A.; Analysis or Interpretation - E.Ö.; Literature Search - S.A.; Writing Manuscript - E.Ö.; Critical Review - E.Ö., K.Ö., S.A.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Morrey BF, Askew LJ, Chao EY. A biomechanical study of normal functional elbow motion. *J Bone Joint Surg Am* 1981; 63: 872-7.
2. de Haan J, Schep NW, Tuinebreijer WE, Patka P, den Hartog D. Simple elbow dislocations: a systematic review of the literature. *Arch Orthop Trauma Surg* 2010; 130: 241-9.
3. Bryce CD, Armstrong AD. Anatomy and biomechanics of the elbow. *Orthop Clin North Am* 2008; 39: 141-54.
4. Mansat P, Morrey BF. The column procedure: a limited lateral approach for extrinsic contracture of the elbow. *J Bone Joint Surg Am* 1998; 80: 1603-15.
5. Chuinard C. The stiff elbow. *Current Orthopaedic Practice* 2008; 19: 602-7.
6. Davila, SA, Johnston-Jones, K.: Managing the stiff elbow: operative, nonoperative, and postoperative techniques. *J Hand Ther* 2006; 19: 268-81.
7. Nandi S, Maschke S, Evans PJ, Lawton JN. The stiff elbow. *Journal of Hand Surgery* 2009; 4: 368-79.
8. Charalambous CP, Morrey BF. Posttraumatic elbow stiffness. *J Bone Joint Surg Am* 2012; 94: 1428-37.
9. Hildebrand KA, Zhang M, Salo PT, Hart DA. Joint capsule mast cells and neuropeptides are increased within four weeks of injury and remain elevated in chronic stages of posttraumatic contractures. *J Orthop Res* 2008; 26: 1313-9.
10. Barron OA, Catalano LW. The stiff elbow: Diagnosis and treatment. *Curr Opin Orthop* 2002; 13: 323-327.
11. Tosun B, Gundes H, Buluc L, Sarlak AY. The use of combined lateral and medial releases in the treatment of post-traumatic contracture of the elbow. *Int Orthop* 2007; 31: 635-8.
12. Garg B, Dua A, Sharma V. Stiff elbow. *Pb Journal of Orthopaedics* 2009; 11.
13. Morrey BF. Post-traumatic contracture of the elbow. Operative treatment including distraction arthroplasty. *J Bone Joint Surg Am* 1990; 72: 601-18.
14. Ball CM, Meunier M, Galatz LM, Calfee R, Yamaguchi K. Arthroscopic treatment of post-traumatic elbow contracture. *J Shoulder Elbow Surg* 2002; 11: 624-9.
15. Tucker SA, Savoie FH, O'Brien MJ. Arthroscopic management of the post-traumatic stiff elbow. *J Shoulder Elbow Surg* 2011; 20: 83-9.
16. Peden JP, Savoie FH. Arthroscopy of the Stiff Elbow. Techniques in orthopaedics. surgical management of complex elbow problems: Update 2006; 21: 331-9.
17. Mansat P, Morrey BF, Hotchkiss RN. Extrinsic contracture: the column procedure, lateral and medial capsular releases. In Morrey BF, ed. *The*

- Elbow and Its Disorders, 3rd ed. Philadelphia: WB Saunders, 2000:447-56.
18. Vingerhoeds B, Degreef I, De Smet L. Debridement arthroplasty for osteoarthritis of the elbow (Outerbridge-Kashiwagi procedure). *Acta Orthop Belg* 2004; 70: 306-10.
19. Morrey BF. Primary Degenerative arthritis of the elbow treatment by ulnohumeral arthroplasty. *J Bone Joint Surg Br* 1992; 74: 409-13.
20. Hertel R, Pisan M, Lambert S, Ballmer F. Operative management of the stiff elbow: sequential arthrolysis based on a transhumeral approach. *J Shoulder Elbow Surg* 1997; 6: 82-8.
21. Wang J, Li H, Zheng Q, Feng G, Li J, Pan Z. Distraction arthrolysis of posttraumatic elbow stiffness with a hinged external fixator. *Orthopedics* 2012; 35: e1625-30.
22. Forthman CL, Jupiter JB. Surgical Approach to the Posttraumatic Stiff Elbow. *Techniques in Shoulder & Elbow Surgery* 2004; 5: 219-230.
23. Lindenhovius AL, Linzel DS, Doornberg JN, Ring DC, Jupiter JB. Comparison of elbow contracture release in elbows with and without heterotopic ossification restriction motion. *J Shoulder Elbow Surg* 2007; 16: 621-5.
24. Liu S, Fan CY, Ruan HJ, Li FF, Tian J. Combination of arthrolysis by lateral and medial approaches and hinged external fixation in the treatment of stiff elbow. *J Trauma* 2011; 70: 373-6.
25. Ehsan A, Huang JI, Lyons M, Hanel DP. Surgical management of posttraumatic elbow arthrofibrosis. *J Trauma Acute Care Surg* 2012; 72: 1399-403.
26. Kayalar M, Ozerkan F, Bal E, Toros T, Ademoğlu Y, Ada S. Elbow arthrolysis in severely stiff elbows. *Arch Orthop Trauma Surg* 2008; 128: 1055-63.
27. Hildebrand KA, Zhang M, van Snellenberg W, King GJ, Hart DA. Myofibroblast numbers are elevated in human elbow capsules after trauma. *Clin Orthop Relat Res* 2004; 189-97.
28. Cohen MS, Schimmel DR, Masuda K, Hastings H, Muehleman C. Structural and biochemical evaluation of the elbow capsule following trauma. *J Shoulder Elbow Surg* 2007; 16: 484-90.